

## RESPONSE OF FABA BEAN (*Vicia faba*, L.) PLANTS TO SEED-TREATING WITH GARLIC EXTRACT, SALICYLIC ACID AND PACLOBUTRAZOL

Wanas, A. L.

Dep. of Agric. Botany, Fac. of Agric., Moshtohor, Benha Univ., Egypt

### ABSTRACT

Application of garlic extract (GE) at 100&200ml/l, salicylic acid (SA) at 100&200 ppm and paclobutrazol (PP<sub>333</sub>) at 10&20 ppm as soaking treatments for faba bean seeds before sowing during 2004/05 and 2005/06 seasons, significantly enhanced many of the vegetative growth characters as plant height, No. of branches, No. of leaves, dry weight of both stems and leaves and total leaf area/plant. Yet, significant reduction in the plant height existed only with PP<sub>333</sub> at its two applied concentrations. Besides, all applied treatments obviously increased photosynthetic pigments, NPK, crude protein and total sugars concentrations in the leaves of treated plants at 75 days after sowing. In addition, different applied treatments positively altered many anatomical features of stems and leaflet blade of treated plants. Since, all applied treatments caused an obvious increase in the thickness of stem wall and its comprising tissues as epidermis, cortex and parenchymatous pith as well as thickness of midrib, lamina, upper and lower epidermis, palisade and spongy tissues in leaves. Also, dimensions of vascular bundles, thickness of phloem and xylem tissues and number of xylem vessels/bundle were increased in both stems and leaves of treated plants. Moreover, with the onset of flowering, different applied seed-soaking treatments significantly increased number of formed flowers and settled pods/plants, whereas reduced the percentages of flowers and pods shedding, in turn significantly increased number of mature (yielded) pods and the final seed yield comparing with those of untreated plants. Furthermore, concentrations of NPK, crude protein, sugars and total carbohydrates in the seeds were also increased as affected by the applied treatments. Hence, it could be recommended the applying of GE, SA and PP<sub>333</sub> as seed-soaking treatments for reducing the abscission of flowers and pods in faba bean plants which consequently reflect upon obvious increase in the final seed yield.

### INTRODUCTION

The phenomena of buds, flowers and immature pod shedding of faba bean usually took place in serious values leading to great reduction in seed yield of this economical plant. Therefore, plant physiologists and breeders are studying intensively the problem of shedding, in order to find out a solution for reducing the high percentage of buds, flowers and immature pods that fail to develop into fully mature pods in this plant. In this respect, many trials have been carried out for increasing flower set, minimizing pre-harvest abscission of immature fruits of faba bean or other plants by the use of different factors including plant growth substances (Abd El-Dayem and El-Deeb, 2000 and Ahmed, 2002), mineral nutrients (Wanas, 2002a) and some natural plant extracts (Atawia and El-Desouky, 1997 and Wanas, 2002b).

Here, garlic extract, salicylic acid and paclobutrazol were used as soaking treatments for faba bean seeds to improve growth and reduce flowers and immature pods shedding of this economical plant.

Wanas, A. L.

from the middle part of the 4<sup>th</sup> apical internode, while those of leaves (1cm<sup>2</sup>) were taken from the middle part of certain leaflet blade of the 4<sup>th</sup> apical leaf on the main stem. The specimens were killed and fixed for at least 48 hours in F.A.A. solution (10 ml formalin + 5 ml glacial acetic acid + 85 ml ethyl alcohol 70%), washed in 50% ethyl alcohol, dehydrated in a series of ethyl alcohols (70, 90, 95 and 100%), infiltrated in xylene, embedded in paraffin wax of a melting point 60-63°C (Sass, 1950), sectioned at 20  $\mu$  using a rotary microtome, double stained with fast green and safranin (Johanson, 1940), cleared in xylene and mounted in Canada balsam.

The prepared sections were microscopically examined. Counts and measurements ( $\mu$ ) were taken using a micrometer eye piece. Averages of readings from 4 sections / treatment were calculated.

#### V-Flowering as well as yield and its components characters:

Ten plants per each treatment were randomly chosen and labeled in the field from the start of flowering to harvesting time and the following characters were studied and recorded:

- No. of opened flowers/plant: Counting was started at 60 days of plant age with 3 days intervals until 100 days.
- No. of setted pods/plant: Counting was started at 75 days of plant age with 3 days intervals until 125 days.
- No. of survived (mature) pods/plant: It was recorded at harvest time.

d) % of flower shedding = 
$$\frac{\text{Total No. of flowers/plant} - \text{No. of setted pods/plant}}{\text{Total No. of flowers/plant}}$$

e) % of pod shedding = 
$$\frac{\text{No. of setted pods/plant} - \text{No. of survived pods/plant}}{\text{Total No. of setted pods/plant}}$$

- No. of seeds/plant, seed yield (g)/plant and seed index [100-seed weight(g)], were recorded at harvesting time.
- Relative total seed yield was calculated as a percentage of control yield.

#### VI-Chemical constituents in the seeds:

Samples from faba bean seeds at harvesting time were taken to determine total nitrogen (Horneck and Miller, 1998), phosphorus (Sandell, 1950) potassium (Horneck and Hanson, 1998), and total and reducing sugars and total carbohydrates (Dubois *et al.*, 1956). Also, crude protein was calculated according to A.O.A.C (1990) using the following equation:  
Crude protein = Total nitrogen X 6.25

#### VII-Statistical analysis:

Data of the vegetative growth, flowering as well as yield and its component were subjected to statistical analysis according to Snedecor and Cochran (1989).

## RESULTS AND DISCUSSION

### 1-Vegetative growth characters:

#### 1-1) Stem characters :

Data in Table (1) indicate that application of garlic extract (GE) at 100&200 ml/l and salicylic acid (SA) at 100&200 ppm as seed-soaking treatments significantly increased the height of treated faba bean plants compared with untreated ones. That was true at the two sampling date during the two seasons. Besides, increases were in parallel to the applied concentrations of GE and SA with superiority of SA in this respect. On the contrary, application of the growth retardant paclobutrazol (PP<sub>333</sub>) at 10&20 ppm caused a significant reduction in this parameter at the two stages of growth during the two growing seasons.

With regard to number of branches and stems dry weight/plant as shown in Table (1), it was found that all applied seed-soaking treatments significantly increased these parameters at the two sampling date in the two grown seasons compared with the control treatment. Here, it could be noticed that PP<sub>333</sub> at 20 ppm gave the highest number of branches/plant followed by GE at 200ml/l, meanwhile GE was the most effective in case of stems dry weight comparing with SA and PP<sub>333</sub>. Increment of branches number/plant is of great importance, because it indicates more dry matter being allocated for the formation of new branches which could be later carried an additional yield.

#### 1-2) Leaf characters :

As shown in Table (1) different estimated growth characters of leaves (number, dry weight and total leaf area/plant) at 75 and 105 days from sowing were significantly increased with all applied treatments compared with those of the control during the two growing seasons. Increases were mostly in parallel to the applied concentrations of GE, SA and PP<sub>333</sub>. Also, the highest increases in leaf parameters, specially at 105 days of plant age existed with GE followed by PP<sub>333</sub> and SA, respectively. Increment of leaf number and total leaf area could be mainly attributed to an increase of the formed branches which, in turn, reflected upon increment of leaves dry weight. In addition, the assimilation rate, i.e., the leaf area in cm<sup>2</sup> required for producing one gram of dry matter, exhibited its significant reduction with all applied treatments. It means that the efficiency of photosynthesis was positively affected by the applied seed-soaking treatments. Also, increases of photosynthetic area and its activity lead to more photosynthates creation and finally could be reflected upon vigorous growth and productivity.

Regarding the enhanceable effect of GE on faba bean growth, it might be due to its stimulatory effect on auxins, gibberelins and cytokinins biosynthesis (Wanas *et al.*, 1998), chlorophyll and carbohydrate formation and protein synthesis (El-Desouky *et al.*, 1998 and Seham, 2002), beside its content of amino acid, sugars, protein, vitamins and other growth factors (Walt and Merrill, 1963).

As for the reducable effect of PP<sub>333</sub> on plant height, Grossman (1990) reported that growth retardants caused shortening in shoots by inhibiting cell division and elongation.

Table (1): Growth behaviour of faba bean plants as affected by garlic extract (GE), salicylic acid (SA) and paclobutrazol (PP<sub>333</sub>) applied as seed-soaking treatments during 2004/05 and 2005/06 seasons

Characters	75										105																	
	Days after sowing										Days after sowing																	
	Plant height (cm)	No. of branches/plant	Stems dry weight(g)/plant	No. of leaves/plant	Leaves dry weight(g)/plant	Total leaf area (cm <sup>2</sup> /plant)	A.R. (cm <sup>2</sup> /g)	Plant height (cm)	No. of branches/plant	Stems dry weight(g)/plant	No. of leaves/plant	Leaves dry weight(g)/plant	Total leaf area (cm <sup>2</sup> /plant)	A.R. (cm <sup>2</sup> /g)	Plant height (cm)	No. of branches/plant	Stems dry weight(g)/plant	No. of leaves/plant	Leaves dry weight(g)/plant	Total leaf area (cm <sup>2</sup> /plant)	A.R. (cm <sup>2</sup> /g)							
Treatments	Season 2004/05																											
Control	74.80	2.00	7.16	35.20	8.24	1768.24	214.35	105.20	2.20	20.91	56.20	16.50	4017.37	243.48	80.00	2.40	9.75	45.60	11.47	2283.18	199.06	113.40	3.00	26.28	71.00	23.74	4894.99	206.18
GE	83.80	3.40	11.48	52.80	13.20	2450.24	185.62	115.40	3.40	29.82	75.20	25.68	5116.99	195.45	81.20	2.00	9.58	40.80	9.88	2100.81	212.63	113.80	2.40	24.38	62.80	19.68	4412.25	224.20
SA	87.80	2.40	11.26	44.20	11.17	2288.97	204.27	118.00	2.80	28.46	69.80	21.38	4748.95	222.10	85.20	2.40	7.95	38.40	9.76	2007.97	205.73	91.60	2.80	24.18	67.80	20.36	4476.96	218.39
PP <sub>333</sub>	59.40	3.80	8.25	48.00	10.34	2089.96	202.12	88.40	3.60	28.69	78.20	24.90	5093.62	204.06	50.40	3.80	8.25	48.00	10.34	2089.96	202.12	88.40	3.60	28.69	78.20	24.90	5093.62	204.06
LSD 0.05	4.12	0.31	0.62	2.21	0.98	78.30	4.66	5.37	0.56	1.83	3.16	2.51	94.21	6.36	4.12	0.31	0.62	2.21	0.98	78.30	4.66	5.37	0.56	1.83	3.16	2.51	94.21	6.36
	Season 2005/06																											
Control	68.00	1.80	6.79	32.80	7.69	1620.01	210.66	95.80	2.00	18.82	52.00	15.01	3786.45	252.26	75.40	2.20	8.44	41.40	10.38	2084.29	200.80	106.20	3.00	24.14	67.20	22.14	4703.35	212.44
GE	79.80	3.00	10.50	48.40	11.79	2314.43	196.30	109.00	3.00	28.42	71.60	23.58	4890.10	207.38	77.60	2.00	9.28	38.20	9.52	1970.36	208.97	102.30	2.00	22.40	56.60	17.44	4110.55	235.70
SA	80.40	2.00	10.66	42.80	10.80	2209.55	204.59	108.20	2.60	27.31	63.60	19.78	4540.54	229.55	80.40	2.80	8.52	44.20	10.20	2063.81	202.33	84.40	3.00	26.28	71.60	21.92	4659.34	212.56
PP <sub>333</sub>	58.00	3.20	8.06	43.60	9.72	1958.37	201.97	81.80	3.40	27.72	75.00	22.72	4772.52	210.08	58.00	3.20	8.06	43.60	9.72	1958.37	201.97	81.80	3.40	27.72	75.00	22.72	4772.52	210.08
LSD 0.05	4.85	0.26	0.76	2.14	0.81	68.00	3.81	6.67	0.49	2.05	2.98	2.18	78.16	7.68	4.85	0.26	0.76	2.14	0.81	68.00	3.81	6.67	0.49	2.05	2.98	2.18	78.16	7.68

\* A.R. = Assimilation rate

The action of PP<sub>333</sub> as a growth retardant has been attributed to its inhibition of GAs biosynthesis (Grossmann, 1990, and Bondok *et al.*, 1995). However, enhancement of branching as a result of PP<sub>333</sub> application might be due to its stimulative effect on cytokinins biosynthesis that had know enhanceable effect on branching (Abd El-Dayem and El-Deeb, 2000). Cytokinins have an important role in stimulating growth and development of lateral buds by increasing their sink capacity and promoting cell division and elongation (Chen, 1997).

With regard to the promotive effect of SA on faba bean growth, it might be due to its stimulatory effect on biosynthesis of the growth promotive hormones, i.e., gibberellins, auxins and cytokinins (Shehata *et al.*, 2000).

#### II-Photosynthetic pigments :

As shown in Table (2), the all applied seed-soaking treatments obviously increased photosynthetic pigments as chlorophyll a, b and carotenoids in the leaves of treated faba bean plants during the two seasons compared with those of untreated ones. Increases were mostly in parallel to the applied concentrations of GE, SA or PP<sub>333</sub>. Besides, the highest increments in chlorophyll a, b as well as total determined pigments were obtained with PP<sub>333</sub> followed by GE, vice versa in case of carotenoids. Meanwhile, SA ranked the last in all cases. In addition, the positive effect of such treatments on photosynthetic pigments may be attributed, in part, to the efficient plant growth and in another to their enhanceable effect on the endogenous cytokinins level (findings of Wanas *et al.*, 1998 for GE, Abd El-Dayem and El-Deeb, 2000 for PP<sub>333</sub> and Shehata *et al.*, 2002 for SA). Cytokinins have been established to induce the biosynthesis of chloroplast pigments in many plants (Fletcher and Arnold, 1988 and Bondok *et al.*, 1995).

Table (2) : Photosynthetic pigments concentrations (mg/g f.wt.) in faba bean leaves as affected by garlic extract (GE), salicylic acid (SA) and paclobutrazol (PP<sub>333</sub>) applied as seed-soaking treatments during 2004/05 and 2005/06 seasons.

Characters Treatments	Chlorophylls				Carotenoids		Total determined pigments		
	a		b		$\bar{X}$	$\pm\%$	$\bar{X}$	$\pm\%$	
	$\bar{X}$	$\pm\%$	$\bar{X}$	$\pm\%$					
Season 2004/05									
Control	0.29	0.00	0.44	0.00	0.56	0.00	1.91	0.00	
GE	100ml	0.95	+8.74	0.43	+9.09	0.86	+13.79	2.09	+9.42
	200ml	1.00	+12.36	0.58	+22.73	0.71	+22.45	2.25	+17.8
SA	100ppm	0.94	+5.62	0.45	+3.27	0.85	+17.07	2.04	+6.81
	200ppm	0.98	+10.11	0.49	+11.36	0.86	+13.79	2.13	+11.52
PP <sub>333</sub>	10ppm	0.98	+10.11	0.52	+18.18	0.82	+6.90	2.12	+10.99
	20ppm	1.08	+21.35	0.58	+31.82	0.86	+13.79	2.32	+21.48
Season 2005/06									
Control	0.78	0.00	0.39	0.00	0.49	0.00	1.68	0.00	
GE	100ml	0.80	+2.56	0.44	+12.82	0.59	+18.37	1.82	+8.64
	200ml	0.91	+16.87	0.49	+23.64	0.81	+24.49	2.01	+21.08
SA	100ppm	0.80	+2.56	0.43	+10.26	0.55	+12.24	1.78	+7.22
	200ppm	0.83	+12.82	0.43	+10.26	0.59	+20.41	1.90	+14.46
PP <sub>333</sub>	10ppm	0.92	+17.95	0.47	+20.51	0.53	+6.18	1.92	+13.66
	20ppm	0.94	+20.51	0.52	+28.21	0.58	+14.29	2.04	+22.69

$\pm\% = \pm\%$  relative to the control value

Table (4): Anatomical features of faba bean stems as affected by garlic extract (GE), salicylic acid (SA) and paclobutrazol (PP<sub>100</sub>) applied as seed-soaking treatments during 2005/06 season

Treatments	Counts and measurements (μ)		Dimension of whole section of hollow pith	Thick. of stem wall	Thick. of epidermis	Thick. of cortex	Thick. of parenchyma in pith	No. of vascular bundles	Length of large bundle	Width of large bundle	Thick. of phloem tissue	Thick. of xylem tissue	No. of Xylem vessels	Thick. of pericortical bundle	Thick. of pericyclic fibers
	X	%													
Control	X	5208.8	2552.0	1328.4	55.8	452.7	819.90	29.0	441.9	466.2	65.7	378.2	50.3	50.3	297.9
	%	6496.7	2818.3	1839.2	59.4	558.5	1221.3	32.0	553.5	539.1	94.5	459.0	67.3	67.3	425.7
GE	X	124.7	110.4	138.5	106.5	123.4	149.0	110.3	125.3	115.6	143.8	122.0	133.8	142.9	142.9
	%	6998.7	3422.2	1739.2	81.2	540.9	1186.2	32.5	626.4	644.4	103.5	540.9	70.5	70.5	387.0
200ppm	X	134.4	134.1	130.9	109.7	110.5	144.7	112.1	141.8	138.2	157.5	143.8	140.2	140.2	128.9
	%	5758.5	2882.1	1447.2	57.6	459.8	919.8	32.0	575.1	609.3	81.0	494.1	59.0	59.0	328.8
100ml/l	X	110.5	112.2	108.9	102.2	103.8	112.2	110.3	130.1	130.7	119.6	131.3	115.3	115.3	109.7
	%	6245.0	2925.8	1659.6	55.8	483.7	1115.1	29.8	594.4	567.0	81.9	512.5	63.3	63.3	358.2
200ml/l	X	119.9	114.6	124.9	100.0	108.0	138.00	102.8	134.5	121.6	124.7	130.2	125.8	125.8	120.2
	%	5749.8	3027.2	1360.8	63.0	468.0	829.8	30.5	470.7	541.8	81.0	389.7	55.8	55.8	327.6
10ppm	X	110.4	118.6	102.4	112.9	103.4	101.2	105.2	106.5	116.2	119.6	103.6	110.9	110.9	110.0
	%	7251.5	3978.9	1636.3	64.8	595.8	975.7	32.5	527.7	715.5	90.0	507.7	65.5	65.5	458.1
PP <sub>100</sub>	X	139.2	155.9	123.2	116.1	131.6	119.0	112.1	135.3	153.5	137.0	135.0	130.2	130.2	153.8
	%	139.2	155.9	123.2	116.1	131.6	119.0	112.1	135.3	153.5	137.0	135.0	130.2	130.2	153.8

\*Control values are considered as 100 %

Stem dimension - hollow pith

Stem wall =




Fig. (1) Transverse sections through the middle part of the 4<sup>th</sup> apical internode of the main stem of faba been as affected by GE, SA and PP<sub>233</sub> applied as seed – soaking treatments (X24)

a) Control

b) Garlic extract (GE) at 200 ml/l

c) Salicylic acid (SA) at 200ppm

d) Paclobutrazol (PP<sub>233</sub>) at 20 ppm

Abb : ep. = epidemis , cx. = cortex, pa. pi. = parenchymatous pith, p.f.= pericyclic fibers, ph. = phloem tissue, xy. = xylem tissue and s.w. = stem wall.

The increase in stem dimension due to GA, SA and PP<sub>333</sub> at the two concentrations used may be attributed to the increase in cortex and parenchymatous pith thickness resulted from their action in cell division and enlargement. Barlow *et al.*, (1991) stated that the increase in stem diameter by uniconazol may be attributed to its effects on promotion of lateral cell division and enlargement. Furthermore, activation in cell division occurred in vascular cambium initial cells forming more xylem and phloem elements.

Herein, of interest to note that these positive responses of different anatomical aspects to GE, SA and PP<sub>333</sub> treatments were completely reflected upon vegetative growth and productivity of treated plants. So, present study revealed those increase of xylem tissues, i.e., the route of mineral nutrients and water translocation from roots to leaves and the phloem tissue, i.e., the pathway of different assimilates from leaves to seed and other sinks. Thereby, improvement of translocation events directly may be considered a direct reason for increment the final seed yield.

#### b) Leaflet blade anatomy :

Data in Table (5) and Fig. (2) indicate that most of the studied anatomical features of faba bean leaflet blades were increased with the assigned treatments of GE, SA and PP<sub>333</sub>. Among these anatomical features were the most important ones, i.e., thickness of midrib region, length and width of vascular bundle, thickness of phloem and xylem tissues and number of xylem vessels in the vascular bundle. Once again GE was the most effective compared with the two growth regulators (SA and PP<sub>333</sub>) regarding the above mentioned characters.

On the other hand, all applied treatments obviously increased thickness of lamina. Also, increment of lamina thickness was accompanied with increases in the thickness of its comprising tissues, i.e., upper and lower epidermis, palisade and spongy tissues. Here, increase values were mostly in parallel to the applied concentrations of GE, SA or PP<sub>333</sub>. Also, GE was the most effective treatment followed by PP<sub>333</sub>, while SA ranked the last one in this concern.

In general, these positive alterations in stem and leaf anatomy of faba bean plants treated with GE, SA and PP<sub>333</sub> as seed-soaking application led to vigorous growth (Table, 1) causing induction of flowering and fruit setting of treated plants (Table, 6). That as will mentioned afterwards reflected upon significant increases in the final seed yield. Similar results have been reported about yield increases through doing alterations in the anatomical features of some economical plants as affected by natural extracts (Wanas *et al.*, 1998 and Wanas, 2002b), and PP<sub>333</sub> (Ahmed, 2002 and Hyam, 2006).



Table (5) : Anatomical features of faba bean leaf blades as affected by garlic extract (GE), salicylic acid (SA) and paclobutrazol (PP<sub>333</sub>) applied as seed-soaking treatments during 2005/06 season

Counts and measurements (a)		Thick of midrib	length of main vascular bundle	Width of main vascular bundle	Thick of phloem tissue	Thick of xylem tissue	No. of xylem vessels/main v. bundle	Thick of lamina	Thick of upper epidermis	Thick of lower epidermis	Thick of palisade tissue	Thick of spongy tissue
Treatments	*Control	X 999.9	272.7	183.6	55.8	216.9	23.5	312.3	36.0	30.6	90.0	155.7
	GE 100ml/l	X 1278.9	330.0	272.7	63.0	267.0	32.5	391.6	36.7	33.3	102.6	207.0
	200ppm	X 1174.5	353.6	260.8	71.1	242.5	35.8	405.9	42.3	34.2	103.7	218.7
		% 117.5	133.3	152.9	127.4	134.9	152.3	130.0	117.5	111.8	120.7	140.5
SA	100ml/l	X 1118.7	318.6	268.0	57.6	261.0	30.3	361.0	38.7	33.3	104.4	185.4
		% 111.9	116.8	150.9	103.2	120.3	128.9	135.9	107.5	108.8	116.0	110.1
	200ml/l	X 1155.6	352.8	286.2	61.2	291.6	31.8	397.8	41.4	34.2	117.0	205.2
		% 115.6	129.4	155.9	109.7	134.4	135.3	127.4	115.0	111.8	130.0	131.8
	10ppm	X 1056.6	359.1	275.4	58.5	300.6	27.8	343.8	36.9	30.6	95.4	180.9
		% 105.7	131.7	150.0	104.8	138.6	118.3	110.1	102.5	100.0	106.0	116.2
PP <sub>333</sub>	20ppm	X 1135.8	362.7	277.2	61.2	301.5	28.5	405.0	41.4	34.2	103.2	223.2
		% 113.6	133.0	151.0	109.7	139.0	121.3	129.7	115.0	111.8	118.0	143.4

\*Control values are considered as 100 %

Wanas, A. L.




Fig. (2) Transverse sections through the leaflet blade of the 4<sup>th</sup> apical leaf on the main stem of faba bean as affected by GE, SA and PP<sub>233</sub> applied as seed - soaking treatments (X60)

a) Control

b) Garlic Extract (GE) at 200 ml/l

c) Salicylic acid (SA) at 200ppm

d) Paclobutrazol (PP<sub>233</sub>) at 20 ppm

Abb : up. ep. = upper epidermis , l. ep. = lower epidermis, pa. = palisade tissue, sp.= spongy tissue, ph. = phloem tissue, xy. = xylem tissue and v.b. = vascular bundle.

**V-Reproductive growth:**

**V-1) Flower formation and shedding :**

Data in Table (6) indicate that different applied seed-soaking treatments (GE, SA, and PP<sub>333</sub> at the two assigned concentrations of each) caused a significant increase in the number of flowers/plant and a significant reduction in the percentage of shedded flowers compared with the control during the two growing seasons. Here, GE was the most effective in this respect followed by PP<sub>333</sub> and SA respectively.

**Table (6) : Flowering as well as yield and its characteristics of faba bean plant as affected by garlic extract (GE), salicylic acid (SA) and paclobutrazol (PP<sub>333</sub>) applied as seed-soaking treatments during 2004/05 and 2005/06 seasons**

Characters	No. of flowers /plant	% of flower shedding	No. of setted pods / plant	% of pod shedding	No. of mature pods / plant	No. of seeds/ plant	Seed Index (g)	Seed yield (g/ plant	Relative seed yield (%)	
Treatments										
<b>Season 2004/05</b>										
Control	109.20	74.91	27.40	39.42	16.60	58.20	58.20	33.87	100.00	
GE	100ml	125.40	71.04	38.60	36.07	23.40	67.80	66.40	45.02	132.92
	200ml	134.40	69.79	40.60	35.47	26.20	73.40	67.16	49.30	145.56
SA	100ppm	118.20	71.57	33.60	38.90	21.20	64.60	62.47	40.38	119.18
	200ppm	128.20	71.76	36.20	36.48	23.00	70.60	64.63	45.63	134.72
PP <sub>333</sub>	10ppm	120.20	70.55	35.40	35.59	22.80	63.80	66.90	42.68	126.01
	20ppm	130.60	69.83	39.40	36.55	25.00	67.20	70.09	47.10	139.06
LSD	0.05	6.11	2.34	2.92	1.76	2.21	3.61	2.03	4.11	-
<b>Season 2005/06</b>										
Control	94.60	72.73	25.80	37.21	16.20	54.60	59.40	32.43	100.00	
GE	100ml	113.00	69.73	34.20	33.92	22.60	62.60	67.69	42.37	130.65
	200ml	120.40	68.60	37.80	33.66	25.00	66.40	67.87	45.07	138.98
SA	100ppm	103.40	70.60	30.40	34.21	20.00	61.80	64.12	39.62	122.17
	200ppm	116.20	70.40	34.40	34.86	22.40	66.20	66.16	43.80	135.06
PP <sub>333</sub>	10ppm	110.40	69.35	33.80	34.32	22.20	59.40	67.96	40.37	124.48
	20ppm	118.20	68.87	36.60	34.24	24.20	64.80	68.52	44.40	136.91
LSD	0.05	3.16	1.85	2.24	1.57	1.75	2.94	2.36	3.28	-

**V-2) Pod yield:**

As shown in Table (6) application of GE, SA and PP<sub>333</sub> significantly increased the number of setted pods/plant, whereas significantly reduced the percentage of shedded ones in proportional to the applied concentrations of each. So, in the two assigned seasons, number of mature pods, i.e., the yielded pods were also significantly increased with all applied treatments. Again, GE was the most pronounced in this respect comparing with the two growth regulators, i.e., PP<sub>333</sub> and SA, respectively.

In addition, significant increases were obtained dominantly in the number of seeds/plant, total seed yield/plant and the calculated relative seed yield as affected by the applied treatments. Here, it could be noticed that the highest increases of total seed yield/plant relative to the control yield (100%)

were obtained with GE at 200 ml/l (45.56 & 38.98%) followed by PP<sub>333</sub> at 20 ppm (39.06 & 36.91%) then SA at 200 ppm (34.72 & 35.06%) during 2004/05 & 2005/06 seasons, respectively. Moreover, seed index, i.e., weight of hundred seeds was also positively responded. Since, its significant increase proportionally existed with the two applied concentrations of GE, SA and PP<sub>333</sub>.

Here, it could be concluded that reduction in shedding percentages of flowers and pods, in turn enhancement of pod setting and development obtained with GE, SA and PP<sub>333</sub> treatments may be due to the enhancable effect of such treatments on total sugars, total protein and mineral concentrations in leaves (Table, 3) as well as their stimulatory effect on the increment of endogenous growth promoters (findings of Wanas, *et al.*, 1998 with GE, Abd El-Dayem and El-Deeb, 2000 with PP<sub>333</sub> and Shehata, 2000 with SA). Auxins and other growth promoters has been shown to effectively retard the process of abscission by prevent structural weakening of the abscission zone. However, once weakening of this specialized layer of cells has begun, auxins and other endogenous growth promoters can effectively block the action of ethylene as well as they can retard the development of senescence in the pulvinal cells on the distal side of the abscission zone in faba bean plant. In addition, auxin prevents synthesis of cellulase seem to be in line with the known action of auxin as an abscission inhibitors (Kozłowski, 1979 and Sakr, 1980). Besides, cytokinins have a direct role in stimulating nutrient mobilization and auxin production (Oosterhuis and Janes, 1997). Hence, all of these advantages positively could reflect on flower initiation, flower development and fruit set and growth as well as inhibiting flowers and young pods shedding and hence increasing the final seed yield.

#### IV-NPK and some bioconstituents in the seeds:

As shown in Table (7) different applied seed-soaking treatments obviously increased NPK, crude protein, total sugars and total carbohydrate concentrations in seeds of treated faba bean plants compared with those of untreated ones. Increases were also in proportional to the applied concentration of GE, SA or PP<sub>333</sub>. In addition, GE at 200 ml/l was the most effective either when compared with its lower one or with the two concentrations of SA & PP<sub>333</sub> regarding different estimated constituents.

In general, the applied seed-soaking treatments improved seed yield of faba bean plants (Table, 6) due to increment of flower formation and the reduction of flowers and pods shedding as well as increasing their ability to accumulate more bioconstituents and NPK (Table, 7). These positive effects of GE, SA and PP<sub>333</sub> treatments upon seed yield and its characters could be considered as a reversion of their effects on the early vigorous growth of faba bean plants, specially that obvious increase in total leaf area (Table, 1) and photosynthetic pigments (Table, 2) and their reflection on increasing the net photosynthesis per unit of leaf area (effects at the source) and increasing the seed weights (Table, 6).

The present study strongly admit the use of GE at 200 ml/l, SA at 200 ppm and PP<sub>333</sub> at 20 ppm as soaking treatments with faba bean seeds for getting the highest seed yield with good quality.

Table (7): NPK and some bioconstituent contents (mg/g d.w.) in faba bean seeds as affected by garlic extract (GE), salicylic acid (SA) and paclobutrazol (PP<sub>333</sub>) applied as seed-soaking treatments during 2004/05 and 2005/06 seasons

Determination Treatments	N	P	K	N + P + K		Crude Protein		Reducing sugars	Non reducing sugars	Total sugars		Total carbohydrates	
				$\bar{x}$	$\pm\%$	$\bar{x}$	$\pm\%$			$\bar{x}$	$\pm\%$	$\bar{x}$	$\pm\%$
Season 2004/05													
Control	41.12	2.96	43.45	87.53	0.00	257.00	0.00	13.38	34.94	48.32	0.00	456.40	0.00
GE	44.26	3.24	48.37	95.87	+9.53	276.63	+7.64	14.34	39.18	52.52	+8.69	485.30	+6.33
	200ml/l												
	46.14	3.98	51.75	101.87	+16.38	289.38	+12.21	15.87	43.73	59.60	+23.34	496.95	+8.89
SA	43.58	3.16	44.17	90.91	+3.86	272.38	+5.98	13.92	37.68	51.60	+6.79	476.45	+4.39
	100ppm												
	45.28	3.70	44.55	93.51	+6.83	292.88	+10.07	15.08	42.14	57.22	+18.42	494.65	+8.38
PP <sub>333</sub>	44.56	3.30	47.82	95.68	+9.31	278.50	+8.37	14.22	41.38	55.60	+15.07	487.90	+6.90
	10ppm												
	46.82	3.78	49.34	99.92	+14.16	292.63	+13.86	15.96	45.26	61.22	+26.70	503.35	+10.28
	20ppm												
Season 2005/06													
Control	39.84	2.85	42.10	84.59	0.00	247.75	0.00	12.25	34.23	46.48	0.00	448.65	0.00
GE	41.18	3.35	47.14	91.67	+8.37	257.38	+3.89	13.50	39.35	52.85	+13.70	473.90	+5.63
	200ml/l												
	44.96	4.22	49.72	98.90	+16.92	281.00	+13.42	15.16	42.12	57.28	+23.24	501.80	+11.85
SA	41.62	3.05	43.62	88.29	+4.37	260.13	+5.00	12.40	36.15	48.55	+4.45	463.85	+3.39
	100ppm												
	43.84	3.54	44.86	91.60	+8.29	274.00	+10.60	13.68	40.12	53.70	+15.53	480.45	+9.32
PP <sub>333</sub>	43.18	3.18	44.58	90.94	+7.51	269.88	+8.93	13.64	40.50	54.14	+18.48	481.20	+7.28
	10ppm												
	46.22	3.96	47.85	98.03	+15.89	288.88	+16.60	15.84	43.62	59.46	+27.93	505.20	+12.60
	20ppm												

$\pm\%$  =  $\pm\%$  relative to the control value

## REFERENCES

- A.O.A.C. (1990). Official Method of Analysis, 15<sup>th</sup> Ed., Association of Official Analytical Chemists, Washington DC, USA.
- Abd El-Dayem, H.M.M and A.E.A El-Deeb (2000): Effects of some growth regulators on growth, yield components and some chemical constituents of cotton plants. *Ann. Agric. Sci., Moshtohor*, 38(2) :907-925.
- Ahmed, H.S.A. (2002) : Botanical studies on growth and flower abortion in cotton (*Gossypium barbadense*) plants. M. Sci. Thesis, Fac. of Agric., Moshtohor, Zagazig Univ.
- Artica, R. N. (1996) : Plant Growth Substances. "Principles and Applications" Chapman and Hall, Thomson Pub., New York, pp. 189-201.
- Atawia, A.A.R. and El-Desouky S.A.(1997): Trials for improving fruit set , yield and fruit quality of Washington navel orange by application of some growth regulators and yeast extract as a natural source of phytohormones. *Ann. Agric. Sci., Moshtohor*, 35 (3): 1613 - 1632.
- Barlow, P.W.; Brian, P. and Parker, J.S. (1991): Cellular growth in roots of a gibberellin deficient mutant of tomato (*Lycopersicon esculentum*, Mill.) and its wild-type. *J. Exp. Bot.*, 42(236):339-351.
- Bondok, M.A., Fouad-Nadia, A. and Felafel, M.S.E. (1995) : The effect of some triazol compounds on the productivity of squash plants. *J. Agric. Sci., Mansoura Univ.*, 20(8): 3677-3690.
- Chen, C. M. (1997) : Cytokinin biosynthesis and interconv. *Physiol. Plant.*, 101: 665-673.
- Derieux, M. ; Kerrest, R. and Y. Montalon (1973): Etude de la surface foliaire et de L'activite photosynthétique chez quelques hybrides de maïs . *Ann. Amélior Plantes*, 23:95-107.
- Dubois, M.; Gilles, K.A.; Hamilton, J.; Rebers, P.A. and Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 28:350-356.
- El-Desouky S.A.; Wanas, A.L. and Khedr, Z.M.A. (1996) : Utilization of some natural plant extracts (of garlic and yeast) as seed-soaked materials to squash (*Cucurbita pepo*, L.) I-Effect on growth, sex expression and fruit yield & quality. *Ann. Agric. Sci Moshtohor, Zagazig Univ.*, 25(2): 839 - 854.
- Fletcher, R.A. and Arnold, V. (1986): Stimulation of cytokinins and chlorophyll synthesis in cucumber cotyledons by triadimefon. *Physiol. Plant.*, 66:197-201.
- Grossmann, K. (1990): Plant growth retardants as tolls in physiological research. *Physiol. Plant.*, 78:640-646.
- Horneck, D.A. and Hanson, D. (1998). Determination of potassium and sodium by Flame Emission Spectrophotometry. In *Handbook of Reference Methods for Plant Analysis* pp. 153-155.

- Horneck, D.A. and Miller, R.O. (1998). Determination of total nitrogen in plant tissue. In Handbook of Reference Methods for Plant Analysis. Kalra Y.P.(Ed.): 75-83.
- Hyam, A. Abd El-Gawad (2006): Growth performance of black cumin (*Nigella saliva*, L.) plants using certain growth conditions. Ph. D. Thesis, . Fac. of Agric., Moshlohor, Benha Univ.
- Kozlowski, T. (1979): Shedding of plant parts. Acad. Press, New York & London.
- Johanson, D.A. (1940) ; Plant Microtechnique. New York and London Mc Grow-Hill Book Co. Inc., pp. 27-154.
- Inskoop, W.P. and Bloom, P.R. (1985): Extraction coefficients of chlorophyll a and b in N,N- dimethylformade and 80% acetone . Plant Physiol., 77 : 493 - 485 .
- Oosterhuis, D.M. and Janes, L.(1997): Effect of three plant growth regulators on the growth and yield of cotton in Arkansas. Proc. FAO-IRCRNC: Joint Meeting of Working Groups 4&3 (Cotton Nutrition & Growth Regulators) 20-23 March, 1995, Cairo- Egypt, pp.235-242.
- Sakr, M. (1980): Physiological studies on factors affecting shedding in broad bean. M. Sci. Thesis, Fac. Agric., Ain Shams Univ.
- Sandell, R. (1950). Colorimetric determination of traces of metal 2<sup>nd</sup> Ed. Interscience Publishers., Inc. New York.
- Sass, J.E. (1950) : Botanical Microtechnique . Iowa State Collage Press, Ames., Iowa, pp 228.
- Seham, M.M. Aly (2002): Effect of some agriculture treatments on yield productivity of squash. Ph. D. Thesis, . Fac. of Agric., Moshlohor, Zagazig Univ.
- Snodcor, G.W. and Cochran, W.G. (1989) : Statistical Method 8<sup>th</sup> Ed., Iowa State Univ. Press .Ames. Iowa, USA.
- Shehata, S.A.M.; Saeed, M.A. and El-Nour, M.S. (2000): Physiological response of cotton plant to the foliar spray with salicylic acid. Ann. Agric. Sci., Cairo, 45(1):1-18.
- Wanas, A. L.; (1996): Botanical studies on some economical plants tolerating salinity. Ph. D. Thesis, Fac. of Agric., Moshlohor, Zagazig Univ.
- Wanas, A. L. (2002a): Response of faba bean (*Vicia faba*, L.) plants to foliar spray with some nutrient elements. Ann. Agric. Sci.. Moshlohor, 40(1) :64-82.
- Wanas, A. L. (2002b): Response of faba bean (*Vicia faba*, L.) plants to seed-soaking application with natural yeast and carrot extracts as phytohormones and other growth factors-rich sources. Ann. Agric. Sci.. Moshlohor, 40(1) :83-102.
- Wanas, A. L.; El-Desouky, S.A. and Khedr, Z.M. (1998): Utilization of some natural plant extracts (of garlic and yeast ) as seed-soaked materials to squash (*Cucubita pepo*, L.) II-Effects on the Histological features and the endogenous hormones. J. Ann. of Agric. Sci. Moshlohor, Zagazig Univ., 35(2): 855-878.

Wanas, A. L.

Wareing, P.E. and P.E. Phillips, I.D.J. (1981): Growth and Differentiation in Plants. 3<sup>rd</sup> Ed., Pergamon Press, Oxford.

Watt, B.K. and Merrill, A.L. (1963): Composition of Foods. US Dept. Agric., Handbook No. 8, 190p.

## استجابة تينات الفول لمعاملة البذور بمستخلص الثوم وحمض الملمسليك والباكلوبيوترزول

احمد لطفي ونسمن

قسم النبات الزراعي - كلية الزراعة بمشهور - جامعة بنها - مصر

أدى استخدام مستخلص الثوم بتركيزي 100 و 200 مللي/لتر وحمض الملمسليك بتركيزي 100 و 200 جزء في المليون والباكلوبيوترزول بتركيزي 10 و 20 جزء في المليون كمعاملات نفع لبذور فصول قبل الزراعة في موسمي 2005/2006. عدد الأفرع - عدد الأوراق - الوزن الجاف لكل من السوق والأوراق وكذلك مساحة الأوراق الكلية/نبات. باستثناء النقص المعنوي في ارتفاع النبات الذي ظهر فقط مع المعاملات المستخدمة إلى زيادة واضحة في تركيز صبغات الكلوروفيل - البيروكسين والفسفور والبوتاسيوم - البروتين الخام - السكريات الكلية بأوراق تينات المعاملة بعد 10 يوم من الزراعة، بالإضافة إلى ذلك فقد سببت معاملات نفع البذور المستخدمة تغيرات إيجابية في العديد من الصفات التشريحية للسوق والأوراق. حيث سببت جميع المعاملات المستخدمة زيادة سمك جدار الساق ومكوناته النسيجية (اللبشر - القشرة - الخنازير البرانشيمي) وكذلك زيادة في سمك منطقة العرق الوسطي والفصل وكل من البشرة العليا والبشرة السفلى والنسيج العمادي والنسيج الاسفنجي في اتصال الأوراق. كما زادت أيضا إمداد الحزم الوعائية وسمك لسبجي اللحاء والخشب وكذلك عدد الأوعية الخشبية/حزمة وعائية في سوق وأوراق النباتات المعاملة. علاوة على ذلك، ومع حلول مرحلة الإزهار سببت جميع المعاملات المستخدمة كمواد نفع للبذور زيادة في عدد الأزهار وعدد القرون المتكونة/نبات في حين سببت نقص في النسجة المثوية لتساقط الأزهار والثمار وبالتالي أدت إلى زيادة في عدد الثمار الناضجة (المستيقية) /نبات وكذلك محصول البذور النهائي/نبات.

وكد ظهرت أعلى قيم لمحصول البذور مع معاملة مستخلص الثوم بتركيز 200 مللي/لتر يليها الباكلوبيوترزول بتركيز 20 جزء في المليون ثم حمض الملمسليك بتركيز 200 جزء من المليون. فضلا عن ذلك، فقد زاد تركيز عناصر البيروكسين والفسفور والبوتاسيوم والبروتين الخام و السكريات والكربوهيدرات الكلية في البذور نتيجة لتأثيرها بالمعاملات المستخدمة.

وبناء على ذلك يمكن التوصية باستخدام مستخلص الثوم وحمض الملمسليك وكذلك الباكلوبيوترزول كمعاملات نفع لبذور الفول بهدف تحسين النمو وتقليل تساقط الأزهار والثمار مما يؤدي إلى زيادة محصول البذور النهائي في هذا النبات الاقتصادي.